arXiv:1207.6804v1 [cond-mat.quant-gas] 29 Jul 2012

Response function and elementary excitations of a Bose-Einstein condensed gas with equal Rashba and Dresselhaus spin-orbit coupling

Giovanni I. Martone[1], Yun Li[1], Lev P. Pitaevskii[1,2], and Sandro Stringari[1]
[1] *Dipartimento di Fisica, Università di Trento and INO-CNR BEC Center, I-38123 Povo, Italy and*
[2] *Kapitza Institute for Physical Problems RAS, Kosygina 2, 119334 Moscow, Russia*

By calculating the dynamic density response function we identify the excitation spectrum of a Bose-Einstein condensate with equal Rashba and Dresselhaus spin-orbit coupling at zero temperature. The propagation of phonons is strongly affected by the coupling and the velocity of sound vanishes when one approaches the second-order phase transition between the spin polarized and the zero-momentum quantum phases. We also point out the emergence of a roton minimum in the excitation spectrum for small values of the Raman coupling, when one approaches the transition to the stripe phase.

PACS numbers: 67.85.De, 03.75.Kk, 03.75.Mn, 05.30.Rt

Synthetic gauge fields are a developing field of research in atomic physics. They have been the object of recent relevant experimental [1–6] and theoretical work [7–15], giving rise to the occurrence of new quantum phases exhibiting unique magnetic features. The elementary excitations of such systems are also expected to exhibit novel features [16–21]. Some of these features have been already the object of experimental measurements [5]. In particular the experiment of [5] has shown that the center of mass oscillation of a harmonically trapped Bose-Einstein condensate can be deeply affected by the coupling with the spin degree of freedom, in agreement with the predictions of theory [20].

The purpose of the present work is to study the elementary excitations and the corresponding behavior of the dynamic structure factor of a spin-orbit coupled Bose-Einstein condensate (BEC) at zero temperature, by direct investigation of the response of the gas to a time dependent perturbation. We explore both the the phonon regime of long wavelengths and the region at higher momentum transfer where the spectrum exhibits novel features including the occurrence of a roton minimum. In ultracold gases the excitation spectrum can be measured via two-photon Bragg spectroscopy [22], so our predictions can be relevant for future experiments on spin-orbit coupled BEC's.

We consider a spin 1/2 interacting Bose gas of N particles enclosed in a volume V, characterized by the Hamiltonian (we set $\hbar = m = 1$)

$$H = \sum_i h_0(i) + \sum_{\alpha,\,\beta} \frac{1}{2} \int d^3\mathbf{r}\, g_{\alpha\beta}\, n_\alpha(\mathbf{r}) n_\beta(\mathbf{r}) \tag{1}$$

where $i = 1, \cdots, N$ is the particle index, while α and β are the spin indexes ($\uparrow, \downarrow = \pm$) characterizing the two spin states. The single particle Hamiltonian h_0 is defined by

$$h_0 = \frac{1}{2}\left[\left(p_x - k_0\sigma_z\right)^2 + p_\perp^2\right] + \frac{\Omega}{2}\sigma_x + \frac{\delta}{2}\sigma_z \tag{2}$$

and is characterized by equal contributions of Rashba [23] and Dresselhaus [24] spin-orbit couplings. Ω is the Raman coupling accounting for the transition between the two spin states, k_0 the momentum transfer of the two Raman lasers and δ is an effective magnetic field providing the energy difference between the two single-particle spin states. The spin up and down density operators entering Eq.(1) are defined by $n_\pm(\mathbf{r}) = (1/2)\sum_i \left(1 \pm \sigma_{z,i}\right)\delta(\mathbf{r}-\mathbf{r}_i)$ while $g_{\alpha\beta} = 4\pi a_{\alpha\beta}$ are the relevant coupling constants in the different spin channels, with $a_{\alpha\beta}$ the corresponding s-wave scattering lengths. Finally σ_k, with $k = x,\ y,\ z$, are the usual 2×2 Pauli matrices. The Hamiltonian (1, 2) has been already implemented experimentally [2, 5] and has been recently employed to describe a variety of non trivial quantum phases in Bose-Einstein condensates [11, 12]. It has the peculiar property of violating both parity and time-reversal symmetry.

In the presence of a spin symmetric Hamiltonian ($g_{\uparrow\uparrow} = g_{\downarrow\downarrow} = g$ and $\delta = 0$), the mean field approach predicts the occurrence of three quantum phases [11, 12], depending on the value of the relevant parameters k_0, Ω, the density n, and the interaction parameters

$$G_1 = n\left(g + g_{\uparrow\downarrow}\right)/4\,; \qquad G_2 = n\left(g - g_{\uparrow\downarrow}\right)/4 \tag{3}$$

These are the stripe phase (also called phase I) where Bose-Einstein condensation (BEC) takes place in a combination of two plane waves with opposite momenta, the single plane wave or spin polarized phase II, where BEC takes place in a single plane wave state with momentum different from zero, and the zero momentum phase III where the condensate has zero momentum. The critical value $\Omega^{(\text{I-II})}$ of the Raman coupling characterizing the first order transition between the stripe and the spin polarized phases can be calculated analytically in the small coupling limit $G_1,\, G_2 \ll k_0^2$, where one finds $\Omega^{(\text{I-II})} = 2k_0^2\sqrt{2\gamma/(1+2\gamma)}$ with $\gamma = G_2/G_1$. The transition between the phases II and III takes instead place at the higher value $\Omega^{(\text{II-III})} = 2\left(k_0^2 - 2G_2\right)$, provided the condition $k_0^2 > 4G_2\left(1+\gamma\right)$ is satisfied, and has a second order nature, being characterized by the divergent behavior of the spin polarizability [20]. If $G_2 < 0$ only the phases II and III are available, the stripe phase being always energetically unfavorable.

In order to calculate the dynamic response of the system we add the time dependent perturbation $V_\lambda = -\lambda e^{i(qx-\omega t)}$ + H.c. to the single particle Hamiltonian (2). The perturbation provides momentum transfer q along the most relevant x-direction. The density response function is then calculated through the usual definition $\chi(q,\omega) = lim_{\lambda\to 0}\delta n/(\lambda e^{-i\omega t})$, where δn are the density fluctuations induced by the external perturbation. In the following we will calculate $\chi(q,\omega)$ by solving the time dependent Gross-Pitaevskii equations. We will restrict the analysis to the phases II and III, where the ground state density is uniform and the wave function of the condensate can be written in the simple form

$$\begin{aligned}\psi = \sqrt{n}\Bigg[&\begin{pmatrix}\cos\theta\\ -\sin\theta\end{pmatrix} e^{ik_1 x}\\ &+\begin{pmatrix}u_\uparrow(x)\\ u_\downarrow(x)\end{pmatrix} e^{-i\omega t} + \begin{pmatrix}v^*_\uparrow(x)\\ v^*_\downarrow(x)\end{pmatrix} e^{i\omega t}\Bigg] e^{-i\mu t}\end{aligned} \tag{4}$$

In the phase II the ground state wave function (first term) is characterized by the condition $\theta = \arccos(k_1/k_0)/2$ and the value $k_1 = k_0\sqrt{1-\Omega^2/[4(k_0^2-2G_2)^2]}$ for the momentum where BEC takes place, corresponding to the value $(n_\uparrow - n_\downarrow)/n = k_1/k_0$ for the spin polarization. In the phase III one instead has $k_1 = 0$, $\theta = \pi/4$ and $n_\uparrow = n_\downarrow$.

By calculating the Bogoliubov amplitudes u and v in the linear (small λ) limit we find the result (near the poles one should replace ω with $\omega + i0$)

$$\chi(q,\omega) = \frac{N\left[-q^2\omega^2 + a_1(q)\omega + a_0(q)\right]}{\omega^4 + b_3(q)\omega^3 + b_2(q)\omega^2 + b_1(q)\omega + b_0(q)} \tag{5}$$

where the values of the various parameters depend on whether one is in Phase II or III. In phase II we find

$$\begin{aligned}
a_0 =\ & q^6/4 - \left[(3+2G_2)\,k_1^2 - 1 + 2G_2\right] q^4\\
& + 4\,(1-2G_2)^2\,k_1^2 q^2\\
a_1 =\ & 4k_1 q^3\\
b_0 =\ & q^8/16 - \left(k_1^2 - G_1 + G_2\right) q^6/2\\
& +\big[k_1^4 - 2\left(3G_1 + 4G_2 + 2G_1G_2 + 2G_2^2\right) k_1^2\\
&\quad + 2G_1\,(1-2G_2)\,\big] q^4\\
& + 4\,(1-2G_2)\left[G_2 k_1^2 + G_1\,(1-2G_2) - 2G_2^2\right] k_1^2 q^2\\
b_1 =\ & k_1 q^5 - 4\left(k_1^2 - 1 - 2G_1 - 2G_2\right) k_1 q^3\\
& - 16G_2\,(1-2G_2)\left(k_1^2 - 1\right) k_1 q\\
b_2 =\ & -q^4/2 + 2\left(3k_1^2 - 2 - G_1 + G_2\right) q^2\\
& + 4\,(1-2G_2)\left(2G_2 k_1^2 - 1\right)\\
b_3 =\ & -4k_1 q
\end{aligned}$$

In Phase III we instead obtain the results

$$\begin{aligned}
a_0 &= q^6/4 + (\Omega - 1 + 2G_2)\,q^4 + \Omega\left[\Omega - 2\,(1-2G_2)\right] q^2\\
b_0 &= q^8/16 + \left[\Omega - 2\,(1 - G_1 - G_2)\right] q^6/4\\
&\qquad + \big[\Omega^2 - 4\,(1 - 2G_1 - G_2)\,\Omega\\
&\qquad + 4\,(1-2G_1)\,(1-2G_2)\,\big] q^4/4\\
&\qquad + 2G_1\Omega\left[\Omega - 2\,(1-2G_2)\right] q^2\\
b_2 &= -q^4/2 - \left[\Omega + 2\,(1 + G_1 + G_2)\right] q^2 - \Omega\,(\Omega + 4G_2)\\
a_1 &= b_1 = b_3 = 0
\end{aligned}$$

where, to simplify the notation, we have expressed the quantities q and k_1 in units of k_0, and the quantities Ω, G_1 and G_2 in units of k_0^2. The above equations include all the relevant information relative to the frequency of the elementary excitations (poles of χ), as well as to the dynamic structure factor given, at $T = 0$, by $S(q,\omega) = \pi^{-1}\mathrm{Im}\chi(q,\omega)$ for $\omega \geq 0$ and $S(q,\omega) = 0$ for negative ω. They also permit to calculate the static response function $\chi(q) \equiv \chi(q,\omega = 0)/N$ yielding, for $q \to 0$, the compressibility χ of the gas. It is worth noticing that the odd terms in ω entering the response function identically vanish in the spin unpolarized phase III, but survive in the spin polarized phase reflecting the lack of parity and time reversal symmetry of the ground state wave function in phase II. The condition $\mathrm{Im}\chi(q,\omega) = -\mathrm{Im}\chi(-q,-\omega)$ characterizing the imaginary part of the response function is always satisfied, but the symmetry relationship $\mathrm{Im}\chi(q,\omega) = \mathrm{Im}\chi(-q,\omega)$ is not ensured in the phase II. Despite this asymmetry and the fact that the compressibility exhibits a jump at the transition, the frequencies of the elementary excitations are always continuous functions of Ω. First results for the excitation spectrum of the Hamiltonian (1)-(2) for small and large values of Ω, far from the transition between the plane wave and single minimum phases, have been recently discussed in [21] using a hydrodynamic approach.

The low frequency excitations at small q (sound waves) can be easily obtained by solving the equation $b_2\omega^2 + b_1\omega + b_0 = 0$. By setting $\omega = cq$ we find, in the phase III, the result

$$c_{\mathrm{III}} = \sqrt{2G_1\left(1 - \frac{2k_0^2}{\Omega + 4G_2}\right)} \tag{6}$$

which explicitly shows the strong reduction of the sound velocity caused by the spin-orbit coupling when one approaches the transition to the spin polarized phase II. At the transition the sound velocity vanishes and the elementary excitations exhibit a different q^2 dependence along the x-direction.

In the plane wave, spin polarized phase II the behavior of the sound velocity exhibits a further interesting feature



FIG. 1: (Color online) Sound velocity as a function of the Raman coupling for the following choice of parameters: $G_1/k_0^2 = 0.2$, $G_2/k_0^2 = 0.05$. The two sound velocities in the Phase II correspond to phonons propagating in the direction parallel ($c_{\rm II}^+$) and antiparallel ($c_{\rm II}^-$) to k_1. The horizontal dashed line corresponds to the value $\sqrt{2G_1} = 0.63\,k_0$ of the sound velocity in the absence of spin-orbit and Raman coupling. The vertical dashed lines indicate the Raman frequencies at which the I-II and II-III phase transitions take place.

caused by the lack of symmetry between q and $-q$

$$c_{\rm II}^{\pm} = \frac{k_1\sqrt{2(k_0^2-2G_2)\left[(k_0^2-2G_1-2G_2)G_2k_1^2+k_0^4G_1\right]}}{k_0^4-2G_2k_1^2} \pm \frac{2G_2k_1\left(k_0^2-k_1^2\right)}{k_0^4-2G_2k_1^2} \qquad (7)$$

where the sign $+(-)$ indicates sound waves propagating parallel (anti-parallel) to k_1. Notice that the asymmetry effect in the sound velocity is caused by the spin interaction term G_2. Setting $G_2 = 0$ one finds $c_{\rm II}^+ = c_{\rm II}^- = \sqrt{2G_1[1-\Omega^2/(4k_0^4)]}$. Also in the phase II the sound velocity vanishes when one approaches the transition to the phase III where $k_1 = 0$. The results for the sound velocity are shown in Fig.1 for a configuration with relatively large G_2 emphasizing the difference between $c_{\rm II}^+$ and $c_{\rm II}^-$. Notice that the sound velocity, in the absence of spin-orbit and Raman coupling, would correspond to the value $c = \sqrt{2G_1}$. This value is asymptotically reached only for very large values of Ω.

The strong quenching effect exhibited by the sound velocity near the II-III phase transition is a remarkable consequence of spin-orbit coupling. This effect is surprising especially in the zero momentum phase III where the compressibility of the gas is unaffected by the coupling and given by $\chi = 1/(2G_1)$. The quenching of the sound velocity is the consequence of the strong coupling between density and spin density in the propagation of sound. It also reflects the fact that, as can be inferred from the analysis of Eq.(5), the phonon modes exhaust the compressibility sum rule $\int_{-\infty}^{+\infty} d\omega S(q,\omega)/\omega$ but, differently from ordinary superfluids, they give a vanishingly small contribution to the f-sum rule $\int_{-\infty}^{+\infty} d\omega S(q,\omega)\omega = Nq^2/2$ when one approaches the transition [25]. This behavior shares important analogies with the quenching of the frequency of the center of mass oscillation, recently explored both experimentally [5] and theoretically [20]. Also the static structure factor $S(q) = \int_0^\infty d\omega S(q,\omega)$ is strongly quenched compared to usual BEC's. This results in an enhancement of the quantum fluctuations of the phase as predicted by the uncertainty principle inequality [26].

The peculiar behavior of the excitation spectrum in the phonon regime can be usefully described also using the hydrodynamic formalism where one writes the spin-up and spin-down components of the order parameter in terms of their modulus and phase [21]. In this case one finds four coupled equations instead of two equations as in usual BEC's. In the phonon regime of large wavelengths and small frequencies one can safely neglect the quantum pressure terms. Furthermore, one finds that the phase difference between the two spin components is blocked ($\varphi_\uparrow = \varphi_\downarrow$) [27]. By requiring the vanishing of the difference $\partial_t\varphi_\uparrow - \partial_t\varphi_\downarrow$ between the equations for the two phases, one then derives the non trivial relationship ($\varphi \equiv \varphi_\uparrow = \varphi_\downarrow$)

$$k_0\nabla_x\delta\varphi - k_0^2 Z\left(-\frac{s}{n}\frac{\delta n}{n} + \frac{\delta s}{n}\right) - 2G_2\frac{\delta s}{n} = 0 \qquad (8)$$

between the phase gradient, the density and the spin fluctuations. In the above equation $s = nk_1/k_0$ is the spin density relative to the equilibrium configuration while $Z = \Omega/(2k_0^2(1-k_1^2/k_0^2)^{3/2})$. Eq.(8) permits to reduce the relevant hydrodynamic equations, i.e. the equation of continuity

$$\partial_t\delta n + \nabla\cdot(n\nabla\delta\varphi) - k_0\nabla_x\left[n\left(-\frac{s}{n}\frac{\delta n}{n} + \frac{\delta s}{n}\right)\right] = 0 \qquad (9)$$

and the equation for the phase

$$\partial_t\delta\varphi + k_1\nabla_x\delta\varphi - k_0^2 Z\frac{s}{n}\left(-\frac{s}{n}\frac{\delta n}{n} + \frac{\delta s}{n}\right) + 2G_1\frac{\delta n}{n} = 0\,, \qquad (10)$$

to a closed set of coupled equations for the density and the phase. It is worth pointing out the crucial changes caused by the spin orbit term in the equation of continuity (9). These changes reflect the fact that the physical current of the system is not simply given by the momentum operator, but is also affected by the spin variable. One can identify the current density operator as $\mathbf{j}(\mathbf{r}) = \mathbf{p}(\mathbf{r}) - k_0\sigma_z(\mathbf{r})\hat{i}$, where $\mathbf{p}(\mathbf{r}) = \sum_i\left[\mathbf{p}_i\,\delta(\mathbf{r}-\mathbf{r}_i) + \text{H.c.}\right]/2$ and $\sigma_z(\mathbf{r}) = \sum_i\sigma_{z,i}\,\delta(\mathbf{r}-\mathbf{r}_i)$ are the momentum and spin density, respectively, and $\hat{i}$ is the unit vector along x. The new current operator satisfies the continuity equation $[H, n(\mathbf{r})] = i\nabla\cdot\mathbf{j}(\mathbf{r})$, with $n(\mathbf{r}) = \sum_{\mathbf{i}}\delta(\mathbf{r}-\mathbf{r_i})$ the density operator. In deriving the equation for the phase we have also identified the expression

$$\mu = 2G_1 + 2G_2 - \frac{k_0^2\Omega^2}{8\left(k_0^2-2G_2\right)^2} \qquad (11)$$

for the chemical potential in phase II whose density dependence is affected by spin-orbit coupling only if $G_2 \neq 0$.

In phase III one has instead $\mu = 2G_1 + (k_0^2 - \Omega)/2$, so that μ is continuous at the transition.

The solutions of the hydrodynamic equations reproduce exactly the results (7) for the sound velocity along the x axis. Concerning the sound velocity along the other directions y and z we find that in phase III its value is not modified with respect to the value $\sqrt{2G_1}$ of usual BEC's, while it is affected in phase II if $G_2 \neq 0$.

The hydrodynamic equations further permit to calculate the relative amplitudes of the density, spin density and momentum density oscillations characterizing the propagation of sound. If $G_2 = 0$ we find the result

$$\frac{\delta s}{\delta n} = \frac{k_1}{k_0} + \frac{1}{\sqrt{k_0^2\chi}}\frac{k_0^2\chi_M}{\sqrt{1+k_0^2\chi_M}} \qquad (12)$$

where $\chi = 1/(2G_1)$ is the compressibility and χ_M is the magnetic polarizability, defined by $\chi_M^{-1} \equiv N\partial^2 E/\partial M^2 = k_0^2(Z-1)$ [20], with E and $M = N_\uparrow - N_\downarrow$ the total energy and magnetization of the system, respectively. Equation (12) shows that, near the transition between phases II and III, where χ_M diverges, the density fluctuations are suppressed and the system behaves like an incompressible fluid, despite the fact that the compressibility χ is finite.

A major usefulness of the hydrodynamic equations is that they can be easily extended to trapped non-uniform configurations. In the simplest $G_2 = 0$ case, corresponding to $G_1 = ng/2$, where the wave vector k_1 is density independent, the 3D hydrodynamic equations can be reduced to the compact form:

$$\partial_t^2 \delta n = g\left[(1-1/Z)\,\nabla_x\left(n\nabla_x \delta n\right) + \nabla_\perp\left(n\nabla_\perp \delta n\right)\right] \qquad (13)$$

with n the Thomas-Fermi density profile given, in the presence of harmonic trapping $V_{\rm ho}(\mathbf{r}) = (\omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2)/2$, by an inverted parabola in both phases II and III: $n(\mathbf{r}) = \left[\mu_0 - V_{\rm ho}(\mathbf{r})\right]/g$, with μ_0 fixed by the normalization condition. One can easily check that all the solutions holding for usual BEC's [28] still hold in the presence of spin-orbit coupling, with the simple replacement of the trapping frequency ω_x with $\omega_x\sqrt{1-1/Z}$. This reproduces the result derived in [20] for the dipole oscillation using a sum rule approach and also shows that the frequency of the other hydrodynamic modes involving a motion of the gas along the x axis will be consequently quenched.



FIG. 2: (Color online) Lower branch of the excitation spectrum in Phase II (a) ($\Omega/k_0^2 = 1.8$ green line, $\Omega/k_0^2 = 1.2$, blue line) and in Phase III (b) ($\Omega/k_0^2 = 2.25$) as a function of the wave vector q. In phase II the spectrum is not symmetric and exhibits a roton minimum for negative q whose energy becomes smaller and smaller as one approaches the transition to the stripe phase at $\Omega/k_0^2 = 0.09$. The dotted line corresponds to the usual Bogoliubov dispersion in the absence of spin-orbit and Raman coupling. The other parameters: $G_1/k_0^2 = 0.12$, $\gamma = G_2/G_1 = 10^{-3}$.

In the last part of this work we discuss the behavior of the excitation spectrum at finite momentum transfer q. The poles of Eq.(5) actually provide two separated branches, the lower one approaching the phonon dispersion at small q. A very peculiar feature of the lower branch is exhibited in the plane wave spin polarized phase II for negative values of q, resulting in the emergence of a roton minimum [21] which becomes more and more pronounced as one approaches the phase transition to the stripe phase I. In Fig.2a we show the excitation spectrum in the phase II, calculated in the experimental conditions of [5]. In Fig.2b we instead show the excitation spectrum in the phase III which exhibits symmetry by inversion of q into $-q$. The physical origin of the roton minimum is quite clear. In the phase II the ground state is degenerate and it is very favorable for atoms to be transferred from the BEC state at $p = k_1$ to the empty state at $p = -k_1$. The occurrence of the roton minimum is also reflected in a strong enhancement in the static response function $\chi(q)$. Notice that $\chi(q)$, differently from $\omega(q)$, is always a symmetric function of q. The occurrence of the roton minimum in the excitation spectrum and the corresponding enhancement of the static response represent a typical tendency of the system towards crystallization. We have investigated in detail the condition for the roton frequency being equal to zero (corresponding to a divergent behavior for $\chi(q)$). A simple analytic expression for the corresponding value of the Raman coupling Ω is obtained in the weak coupling limit $G_1, G_2 \ll k_0^2$ were we find that the critical value exactly coincides with the value $\Omega_{\text{I-II}} = 2k_0^2\sqrt{2\gamma/(1+2\gamma)}$ characterizing the transition between the plane wave and the stripe phases. For larger values of the coupling constants G_1 and G_2 we expect that the critical value takes place for values of the Raman coupling smaller than the value at the transition, exhibiting the typical spinoidal behavior characterizing the usual first-order liquid-crystal phase transition.

In conclusion we have investigated the dynamic behavior of a Bose-Einstein condensate with spin-orbit coupling, pointing out the occurrence of novel features of high relevance for future experiments, like the strong quenching exhibited by the sound velocity near the second-order transition the plane wave and the single minimum phases, and the occurrence of a roton minimum in the excitation spectrum in the spin polarized plane wave phase.

This work has been supported by ERC through the QGBE grant.

[1] Y.-J. Lin, R. L. Compton, A. R. Perry, W. D. Phillips, J. V. Porto, and I. B. Spielman, Phys. Rev. Lett. **102**, 130401 (2009); Y.-J. Lin, R. L. Compton, K. Jiménez-García, J. V. Porto, and I. B. Spielman, Nature **462**, 628 (2009); Y.-J. Lin, R. L. Compton, K. Jiménez-García, W. D. Phillips, J. V. Porto, and I. B. Spielman, Nature Phys. **7**, 531 (2011).

[2] Y.-J. Lin, K. Jiménez-García, and I. B. Spielman, Nature **471**, 83 (2011).

[3] M. Aidelsburger, M. Atala, S. Nascimbène, S. Trotzky, Y.-A. Chen, and I. Bloch, Phys. Rev. Lett. **107**, 255301 (2011); P. Hauke *et al.*, arXiv:1205.1398.

[4] L. J. LeBlanc, K. Jiménez-García, R. A. Williams, M. C. Beeler, A. R. Perry, W. D. Phillips, and I. B. Spielman, Proc. Natl. Acad. Sci. USA **109**, 10811 (2012).

[5] S. Chen, J.-Y. Zhang, S.-C. Ji, Z. Chen, L. Zhang, Z.-D. Du, Y. Deng, H. Zhai, and J.-W. Pan, arXiv:1201.6018.

[6] P. Wang, Z.-Q. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai, and J. Zhang, arXiv:1204.1887; L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwierlein, arXiv:1205.3483.

[7] J. Dalibard, F. Gerbier, G. Juzeliūnas, and P. Öhberg, Rev. Mod. Phys. **83**, 1523 (2010).

[8] X.-J. Liu, M. F. Borunda, and X. Liu, and J. Sinova, Phys. Rev. Lett. **102**, 046402 (2009).

[9] T. D. Stanescu, B. Anderson, and V. Galitski, Phys. Rev. A **78**, 023616 (2008); C. Wang, C. Gao, C.-M. Jian, and H. Zhai, Phys. Rev. Lett. **105**, 160403 (2010); C.-J. Wu, I. Mondragon-Shem and X.-F. Zhou, Chin. Phys. Lett. **28**, 097102 (2011).

[10] S. Sinha, R. Nath, and L. Santos, Phys. Rev. Lett. **107**, 270401 (2011); H. Hu, B. Ramachandhran, H. Pu, and X.-J. Liu, Phys. Rev. Lett. **108**, 010402 (2012).

[11] T.-L. Ho and S. Zhang, Phys. Rev. Lett. **107**, 150403 (2011).

[12] Y. Li, L. P. Pitaevskii, and S. Stringari, Phys. Rev. Lett. **108**, 225301 (2012).

[13] T. Ozawa and G. Baym, Phys. Rev. A **85**, 013612 (2012); T. Ozawa and G. Baym, Phys. Rev. Lett. **109**, 025301 (2012).

[14] J. P. Vyasanakere and V. B. Shenoy, Phys. Rev. B **83**, 094515 (2011); J. P. Vyasanakere, S. Zhang, and V. B. Shenoy, Phys. Rev. B **84**, 014512 (2011).

[15] M. Gong, S. Tewari, and C. Zhang, Phys. Rev. Lett. **107** 195303 (2011); H. Hu, L. Jiang, X.-J. Liu, and H. Pu, Phys. Rev. Lett. **107**, 195304 (2011); Z.-Q. Yu and H. Zhai, Phys. Rev. Lett. **107**, 195305 (2011).

[16] E. van der Bijl and R. A. Duine, Phys. Rev. Lett. **107**, 195302 (2011).

[17] Y. Zhang, L. Mao, and C. Zhang, Phys. Rev. Lett. **108** 035302 (2012); Y. Zhang, G. Chen, and C. Zhang, arXiv:1111.4778.

[18] B. Ramachandhran, B. Opanchuk, X.-J. Liu, P. Han, P. D. Drummond, and H. Hu, arXiv:1201.1471.

[19] Z. Chen and H. Zhai, arXiv:1204.5121.

[20] Y. Li, G. I. Martone, and S. Stringari, arXiv:1205.6398.

[21] W. Zheng, and Z. Li, Phys. Rev. A **85** 053607 (2012).

[22] J. Stenger, S. Inouye, A. P. Chikkatur, D. M. Stamper-Kurn, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **82**, 4569 (1999); D. M. Stamper-Kurn, A. P. Chikkatur, A. Görlitz, S. Inouye, S. Gupta, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. **83**, 2876 (1999); J. Steinhauer, R. Ozeri, N. Katz, and N. Davidson, Phys. Rev. Lett. **88**, 120407 (2002).

[23] Y. A. Bychkov and E. I. Rashba, J. Phys. C **17**, 6039 (1984).

[24] G. Dresselhaus, Phys. Rev. **100**, 580 (1955).

[25] By taking the large ω limit one immediately verifies that the response function (5) is consistent with the f-sum rule. However, when one approaches the II-III phase transition, most of the contribution to this sum rule, in the small q limit, comes form the upper branch of the excitation spectrum.

[26] L. Pitaevskii and S. Stringari, J. Low Temp. Phys. **85**, 377 (1991); Phys. Rev. B **47**, 10915 (1993).

[27] The blocking of the relative phase $\varphi_\uparrow - \varphi_\downarrow$ is the consequence of the equation $\partial s/\partial t$ for the spin density which imposes that the quantity $(\varphi_\uparrow - \varphi_\downarrow)$ be small if $\Omega \gg \omega$. The oscillation of the relative phase plays a crucial role in the excitation of the upper branch, whose frequency is of the order of Ω.

[28] S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).